

Heat Shock Protein 27 (HSP27) Ab-1 (Clone G3.1)

Mouse Monoclonal Antibody

Cat. #MS-101-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Purified Ab with BSA and Azide)**Cat. #MS-101-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml)** (Purified Ab without BSA and Azide)**Cat. #MS-101-B0, -B1, or -B (0.1ml, 0.5ml, or 1.0ml at 200µg/ml)** (Biotin-Labeled Ab with BSA and Azide)**Cat. #MS-101-R7 (7.0ml)** (Ready-to-Use for Immunohistochemistry)**Cat. #MS-101-PCS (5 Slides)** (Positive Control for Histology)**Cat. #MS-101-PCL (0.1ml)** (Positive Control for Western Blot)**Please note this data sheet has been changed effective December 6, 2011**

Description: Hsp27 is identical to the estrogen-induced "p29" and "24K" protein. About 50% of breast carcinomas are positive for hsp27 especially those that are also positive for estrogen and/or progesterone receptor. HSP27 has also been implicated in drug resistance in cancer cells.

Mol. Wt. of Antigen: 24-27kDa**Epitope:** Not determined**Species Reactivity:** Human, Chimpanzee, and Monkey. Does not react with mouse and rat. Others-not known.**Clone Designation:** G3.1**Ig Isotype:** IgG₁**Immunogen:** Partially purified hsp27 (earlier called 24K) protein from breast cancer MCF-7 cells.**Applications and Suggested Dilutions:**

- Western Blotting (Ab 1-2µg/ml for 2hrs at RT)
- Immunohistology (Formalin/paraffin) (Ab 1-2µg/ml for 30 min at RT)
- * [Staining of formalin-fixed tissues REQUIRES boiling tissue sections in 10mM citrate buffer, pH 6.0, (Lab Vision Cat. #AP-9003), for 10-20 min followed by cooling at RT for 20 min.]

The optimal dilution for a specific application should be determined by the investigator.

Positive Control: BT474 or MDA-MB-453 cells. Breast carcinoma.**Cellular Localization:** Cytoplasmic / nuclear**Storage and Stability:**

Ab with sodium azide is stable for 24 months when stored at 2-8°C. Antibody WITHOUT sodium azide is stable for 36 months when stored at below 0°C.

Supplied As:

200µg/ml of antibody purified from ascites fluid by Protein G chromatography. Prepared in 10mM PBS, pH 7.4, with 0.2% BSA and 0.09% sodium azide. Also available without BSA and azide at 1mg/ml. Or Prediluted antibody which is ready-to-use for staining of formalin-fixed, paraffin-embedded tissues.

Key References:

1. Edwards DP *et al.* Biochem Biophys Research Commun, 93:804-812, 1980.
2. Ciocca DR *et al.* Breast Cancer Research and Treatment, 20:33-42, 1991.

Limitations and Warranty:

Our products are intended FOR RESEARCH USE ONLY and are not approved for clinical diagnosis, drug use or therapeutic procedures. No products are to be construed as a recommendation for use in violation of any patents. Lab Vision makes no representations, warranties or assurances as to the accuracy or completeness of information provided on our data sheets and website. Our warranty is limited to the price paid for the product. Lab Vision is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.

Material Safety Data:

This product is not licensed or approved for administration to humans or to animals other than the experimental animals. Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. Appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. The material contains 0.09% sodium azide as a preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material as indicated above. The National Institute of Occupational Safety and Health has issued a bulletin citing the potential explosion hazard due to the reaction of sodium azide with copper, lead, brass, or solder in the plumbing systems. Sodium azide forms hydrazoic acid in acidic conditions and should be discarded in a large volume of running water to avoid deposits forming in metal drainage pipes.

For Research Use Only

Heat Shock Protein 27 (HSP27) Ab-1 (Clone G3.1)

Mouse Monoclonal Antibody

Cat. #MS-101-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Purified Ab with BSA and Azide)**Cat. #MS-101-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml)** (Purified Ab without BSA and Azide)**Cat. #MS-101-B0, -B1, or -B (0.1ml, 0.5ml, or 1.0ml at 200µg/ml)** (Biotin-Labeled Ab with BSA and Azide)**Cat. #MS-101-R7 (7.0ml)** (Ready-to-Use for Immunohistochemistry)**Cat. #MS-101-PCS (5 Slides)** (Positive Control for Histology)**Cat. #MS-101-PCL (0.1ml)** (Positive Control for Western Blot)

Please note this data sheet has been changed effective December 6, 2011

Additional Key References:

1. Berra A; Dutt JE; Nouri M; Foster CS. Heat shock protein expression in human conjunctiva. *Investigative Ophthalmology and Visual Science*, 1994, 35(2):352-7.
2. Kato H; Liu Y; Kogure K; Kato K. Induction of 27-kDa heat shock protein following cerebral ischemia in a rat model of ischemic tolerance. *Brain Research*, 1994, 634(2):235-44.
3. Mehlen P; Arrigo AP. The serum-induced phosphorylation of mammalian hsp27 correlates with changes in its intracellular localization and levels of oligomerization. *Eur J of Biochemistry*, 1994, 221:327-34.
4. Tetu B; Brisson J. Prognostic significance of HER-2/neu oncoprotein expression in node-positive breast cancer. The influence of the pattern of immunostaining and adjuvant therapy. *Cancer*, 1994, 73(9):2359-65.
5. Ungar DR; Hailat N; Strahler JR; Kuick RD; Brodeur GM; Seeger RC; Reynolds CP; Hanash SM. Hsp27 expression in neuroblastoma: correlation with disease stage. *J of the National Cancer Institute*, 1994, 86:780-4.
6. Zhu Y; Tassi L; Lane W; Mendelsohn ME. Specific binding of the transglutaminase, platelet factor XIII, to HSP27. *J of Biological Chemistry*, 1994, 269:22379-84.
7. Elledge RM; Ciocca DR; Langone G; McGuire WL. Estrogen receptor, progesterone receptor, and HER-2/neu protein in breast cancers from pregnant patients. *Cancer*, 1993, 71(8):2499-506.
8. Faucher C; Capdevielle J; Canal I; Ferrara P; Mazarguil H; McGuire WL; Darbon JM. The 28-kDa protein whose phosphorylation is induced by protein kinase C activators in MCF-7 cells belongs to the family of low molecular mass heat shock proteins and is the estrogen-regulated 24-kDa protein. *Journal of Biological Chemistry*, 1993, 268(20):15168-73.
9. Head MW; Corbin E; Goldman JE. Overexpression and abnormal modification of the stress proteins alpha B-crystallin and HSP27 in Alexander disease. *American Journal of Pathology*, 1993, 143(6):1743-53.
10. Hurlimann J; Gebhard S; Gomez F. Oestrogen receptor, progesterone receptor, pS2, ERD5, HSP27 and cathepsin D in invasive ductal breast carcinomas. *Histopathology*, 1993, 23(3):239-48.
11. Iwaki T; Iwaki A; Tateishi J; Sakaki Y; Goldman JE. Alpha B-crystallin and 27-kd heat shock protein are regulated by stress conditions in the central nervous system and accumulate in Rosenthal fibers. *American Journal of Pathology*, 1993, 143(2):487-95.
12. Jaattela M; Wissing D. Heat-shock proteins protect cells from monocyte cytotoxicity: possible mechanism of self-protection. *J of Experimental Med*, 1993, 177:231-6.
13. Lavoie JN; Gingras-Breton G; Tanguay RM; Landry J. Induction of Chinese hamster HSP27 gene expression in mouse cells confers resistance to heat shock. HSP27 stabilization of the microfilament organization. *Journal of Biological Chemistry*, 1993, 268(5):3420-9.
14. Mehlen P; Briolay J; Smith L; Diaz-latoud C; Fabre N; Pauli D; Arrigo AP. Analysis of the resistance to heat and hydrogen peroxide stresses in COS cells transiently expressing wild type or deletion mutants of the Drosophila 27-kDa heat-shock protein. *European Journal of Biochemistry*, 1993, 215(2):277-84.
15. Oesterrich S, Wen C-N, Qiu M, et. al. Small heat shock protein hsp27 is correlated with growth and drug resistance in human breast cancer cell lines. *Cancer Res*, 1993, 53: 4443-8.
16. Yokoyama N; Iwaki T; Goldman JE; Tateishi J; Fukui M. Small heat-shock protein is expressed in meningiomas and in granulofilamentous inclusion bodies. *Acta Neuropathologica*, 1993, 85(3):248-55.
17. Ciocca DR; Lo Castro G; Alonio LV; Cobo MF; Lotfi H; Teyssie A. Effect of human papillomavirus infection on estrogen receptor and heat shock protein hsp27 phenotype in human cervix and vagina. *International Journal of Gynecological Pathology*, 1992, 11(2):113-21.
18. Navarro D; Cabrera JJ; Leon L; Chirino R; Fernandez L; Lopez A; Rivero JF; Fernandez P; Falcon O; Jimenez P; et al. Endometrial stromal sarcoma expression of estrogen receptors, progesterone receptors and estrogen-induced spr27 (24K) suggests hormone responsiveness. *Journal of Steroid Biochemistry and Molecular Biology*, 1992, 41(3-8):589-96.
19. Rizk B; Manners CV; Davies MC; Steer CV; Bell SC; Dillon D; Edwards RG. Immunohistochemical expression of endometrial proteins and pregnancy outcome in frozen embryo replacement cycles. *Human Reproduction*, 1992, 7(3):413-7.
20. Yufu Y; Nishimura J; Nawata H. High constitutive expression of heat shock protein 90 alpha in human acute leukemia cells. *Leukemia Research*, 1992, 16:597-605.
21. Ciocca DR; Jorge AD; Jorge O; Milutin C; Hosokawa R; Diaz Lestren M; Muzzio E; Schulkin S; Schirbu R. Estrogen receptors, progesterone receptors and heat-shock 27-kD protein in liver biopsy specimens from



Heat Shock Protein 27 (HSP27) Ab-1 (Clone G3.1)

Mouse Monoclonal Antibody

Cat. #MS-101-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Purified Ab with BSA and Azide)

Cat. #MS-101-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml) (Purified Ab without BSA and Azide)

Cat. #MS-101-B0, -B1, or -B (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Biotin-Labeled Ab with BSA and Azide)

Cat. #MS-101-R7 (7.0ml) (Ready-to-Use for Immunohistochemistry)

Cat. #MS-101-PCS (5 Slides) (Positive Control for Histology)

Cat. #MS-101-PCL (0.1ml) (Positive Control for Western Blot)

Please note this data sheet has been changed effective December 6, 2011

patients with hepatitis B virus infection. *Hepatology*, 1991, 13(5):838-44.

22. Ciocca DR; Luque EH. Immunological evidence for the identity between the hsp27 estrogen-regulated heat shock protein and the p29 estrogen receptor-associated protein in breast and endometrial cancer. *Breast Cancer Research and Treatment*, 1991, 20(1):33-42.

23. Engel K; Knauf U; Gaestel M. Generation of antibodies against human hsp27 and murine hsp25 by immunization with a chimeric small heat shock protein. *Biomedica Biochimica Acta*, 1991, 50(9):1065-71.

24. Mendelsohn ME; Zhu Y; O'Neill S. The 29-kDa proteins phosphorylated in thrombin-activated human platelets are forms of the estrogen receptor-related 27-kDa heat shock protein. *Proceedings of the National Academy of Sciences of the United States of America*, 1991, 88(24):11212-6.

25. Thor A, Benz C, Moore D, et. al. Stress response protein (srp-27) determination on primary human breast carcinomas: clinical, histologic, and prognostic correlations. *J National Cancer Inst*, 1991, 83:170-8.

26. Courtney SP; Williams S; Mansel RE. Evaluation of immunohistochemical staining of breast tissue by an oestrogen-regulated protein, 24K according to epidemiological and radiological breast cancer risk criteria. *Cancer Letters*, 1990, 54:75-80.

27. Manners CV. Endometrial assessment in a group of infertile women on stimulated cycles for IVF: immunohistochemical findings. *Human Reproduction*, 1990, 5(2):128-32.

28. Pauli D; Tonka CH; Tissieres A; Arrigo AP. Tissue-specific expression of the heat shock protein HSP27 during *Drosophila melanogaster* development. *Journal of Cell Biology*, 1990, 111(3):817-28.

29. Fuqua SA; Blum-Salinger M; McGuire WL. Induction of the estrogen-regulated "24K" protein by heat shock. *Cancer Research*, 1989, 49(15):4126-9.

30. Zantema A; de Jong E; Lardenoije R; van der Eb AJ. The expression of heat shock protein hsp27 and a complexed 22-kilodalton protein is inversely correlated with oncogenicity of adenovirus-transformed cells. *Journal of Virology*, 1989, 63(8):3368-75.

31. Moretti-Rojas I; Fuqua SA; Montgomery RA 3d; McGuire WL. A cDNA for the estradiol-regulated 24K protein: control of mRNA levels in MCF-7 cells. *Breast Cancer Research and Treatment*, 1988, 11(2):155-63.

32. Dressler LG; Ramzy I; Sledge GW; McGuire WL. A new marker of maturation in the cervix: the estrogen-regulated 24K protein. *Obstetrics and Gynecology*, 1986, 68(6):825-31.

33. McGuire WL; Dressler LG; Sledge GW Jr; Ramzy I; Ciocca DR. An estrogen-regulated protein in normal and malignant endometrium. *Journal of Steroid Biochemistry*, 1986, 24(1):155-9.

34. Adams DJ; McGuire WL. Quantitative enzyme-linked immunosorbent assay for the estrogen-regulated M_r 24,000 protein in human breast tumors: Correlation with estrogen and progesterone receptors. *Cancer Research*, 1985, 45:2445-9.

35. Ciocca DR; Puy LA; Edwards DP; Adams DJ; McGuire WL. The presence of an estrogen-regulated protein detected by monoclonal antibody in abnormal human endometrium. *Journal of Clinical Endocrinology and Metabolism*, 1985, 60(1):137-43.

36. Sledge GW Jr; Ramzy I; Dressler LG; Dorr FA; Adams DJ; McGuire WL. Presence of an estrogen-regulated protein in endometrial cancer. *Obstetrics and Gynecology*, 1985, 66(3):423-7.

37. Ciocca DR; Adams DJ; Edwards DP; Bjercke RJ; McGuire WL. Estrogen induced 24K-protein in MCF-7 breast cancer cells is localized in granules. *Breast Cancer Research And Treatment*, 1984, 4:261-8.

38. Adams DJ; Hajj H; Bitar KG; Edwards DP; McGuire WL. Purification of an estrogen-regulated breast cancer protein by monoclonal antibody affinity chromatography. *Endocrinology*, 1983, 113:415-7.

39. Adams DJ; Hajj H; Edwards DP; Bjercke RJ; McGuire WL. Detection of a M_r 24,000 estrogen-regulated protein in human breast cancer by monoclonal antibodies. *Cancer Research*, 1983, 43:4297-301.

40. Ciocca DR; Adams DJ; Edwards DP; Bjercke RJ; McGuire WL. Distribution of an estrogen-induced protein with a molecular weight of 24,000 in normal and malignant human tissues and cells. *Cancer Research*, 1983,43:1204-10.

41. Ciocca DR; Asch RH; Adams DJ; McGuire WL. Evidence for modulation of a 24K protein in human endometrium during the menstrual cycle. *Journal of Clinical Endocrinol and Metabolism*, 1983, 57:496-9.

42. Ciocca DR; Adams DJ; Edwards DP; Bjercke RJ; McGuire WL. Immunohistochemical detection of an estrogen regulated protein by monoclonal antibodies. *Cancer Research*, 1982, 42:4256-8.

