

CD56 / NCAM-1 Ab-4 (56C04; same as 123A8)**Mouse Monoclonal Antibody****Cat. #MS-1149-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml)** (Purified with BSA and Azide)**Cat. #MS-1149-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml)** (Purified without BSA and Azide)**Cat. #MS-1149-R7 (7.0ml)** (Ready-to-Use for Immunohistochemical Staining)**Cat. #MS-1149-PCS (5 Slides)** (Positive Control for Histology)

Please note this data sheet has been changed effective December 9, 2011

Description: Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135kDa isoform is the basic molecule which is glycosylated or sialylated to produce the mature species. NCAM (CD56) is reported to express on most neuroectodermal derived cell lines, tissues, and neoplasms such as retinoblastoma, medullblastoma, astrocytoma, and neuroblastoma. It is also expressed on some mesodermally derived tumors such as rhabdomyosarcoma and also on natural killer cells.

Mol. Wt. of Antigen: 180, 145 and 125kDa**Epitope:** Not determined**Species Reactivity:** Human. Others-not known.**Clone Designation:** 56C04 (same as 123A8)**Ig Isotype / Light Chain:** IgG₁ / κ**Immunogen:** Membrane preparation of a small cell lung carcinoma**Applications and Suggested Dilutions:**

- Immunohistology (Formalin/paraffin)
(Use Ab at 1-2µg/ml for 30 minute at RT)
- * (Staining of formalin/paraffin tissues REQUIRES boiling tissue sections in 10mM citrate buffer, pH 6.0, (**NEOMARKERS'** Cat. #AP9003) for 10-20 min followed by cooling at RT for 20 min.)

The optimal dilution for a specific application under a given set of experimental conditions should be determined by the investigator.

Positive Control: Neuroblastoma.**Cellular Localization:** Cell membrane**Storage and Stability:** Ab with sodium azide is stable for 24 months when stored at 2-8°C. Antibody WITHOUT sodium azide is stable for 36 months when stored at below 0°C.**Supplied As:**

200µg/ml of antibody purified from ascites fluid by Protein G chromatography. Prepared in 10mM PBS, pH 7.4, with 0.2% BSA and 0.09% sodium azide. Also available without BSA and azide at 1mg/ml.

or

Prediluted antibody which is ready-to-use for staining of formalin-fixed, paraffin-embedded tissues.

Key References:

1. Schol DJ; et al. Int J Cancer. Supplement, 1988, 2:34-40

Limitations and Warranty:

Our products are intended FOR RESEARCH USE ONLY and are not approved for clinical diagnosis, drug use or therapeutic procedures. No products are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our data sheets and website. Our warranty is limited to the actual price paid for the product. NeoMarkers is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.

Material Safety Data:

This product is not licensed or approved for administration to humans or to animals other than the experimental animals. Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. Appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. The material contains 0.09% sodium azide as a preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material as indicated above. The National Institute of Occupational Safety and Health has issued a bulletin citing the potential explosion hazard due to the reaction of sodium azide with copper, lead, brass, or solder in the plumbing systems. Sodium azide forms hydrazoic acid in acidic conditions and should be discarded in a large volume of running water to avoid deposits forming in metal drainage pipes.



CD56 / NCAM-1 Ab-4 (56C04; same as 123A8)

Mouse Monoclonal Antibody

Cat. #MS-1149-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Purified with BSA and Azide)

Cat. #MS-1149-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml) (Purified without BSA and Azide)

Cat. #MS-1149-R7 (7.0ml) (Ready-to-Use for Immunohistochemical Staining)


Cat. #MS-1149-PCS (5 Slides) (Positive Control for Histology)

Please note this data sheet has been changed effective December 9, 2011

For Research Use Only



Lab Vision Corporation
46360 Fremont Blvd.
Fremont, CA 94538-6406, USA
US Toll Free: 1 (800) 522-7270
Phone: +1 (269) 544-5600
Fax: 1 (269) 372-2674
www.thermoscientific.com/labvision



Thermo Fisher Scientific
Anatomical Pathology
Tudor Road, Manor Park
Runcorn, Cheshire WA7 1TA, UK
Tel: +44 (0) 1928 534 050
Fax: +44 (0) 1928 534 049
sales.ap.uk@thermofisher.com