

**CD74 Ab-1 (Clone LN2)**

Mouse Monoclonal Antibody

**Cat. #MS-131-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml)** (Purified Ab with BSA and Azide)**Cat. #MS-131-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml)** (Purified Ab without BSA and Azide)**Cat. #MS-131-R7 (7.0ml)** (Ready-to-Use for Immunohistochemistry)**Cat. #MS-131-PCS (5 Slides)** (Positive Control for Histology)**Cat. #MS-131-PCL (0.1ml)** (Positive Control for Western Blot)**Please note this data sheet has been changed effective December 6, 2011**

**Description:** Expression of the invariant chain of the HLA-DR antigen (CD74) by B-lymphocytes begins in the pre-B stage but lost before the final differentiation to plasma cells.

**Comments:** Ab-1 has been shown to be useful in the differential identification in non-Hodgkin's lymphomas of the B-lymphocyte phenotype. It recognizes B-lymphocytes and many B-cell lymphomas.

**Mol. Wt. of Antigen:** 35kDa

**Epitope:** Not determined

**Species Reactivity:** Human and Baboon. Does not react with Rat. Others-not known.

**Clone Designation:** LN2 (Workshop IV)

**Ig Isotype / Light Chain:** IgG<sub>1</sub> / κ

**Immunogen:** SU-DHL-4 lymphoma cells.

**Applications and Suggested Dilutions:**

- Flow Cytometry
- Immunofluorescence
- Western Blotting (Ab 1-2µg/ml for 2hrs at RT)
- Immunohistology (Formalin/paraffin) (Ab 2-4 µg/ml for 30 min at RT)
- \* [Staining of formalin-fixed tissues REQUIRES boiling tissue sections in 10mM citrate buffer, pH 6.0, (Lab Vision Cat. #AP-9003), for 10-20 min followed by cooling at RT for 20 min.]

The optimal dilution for a specific application should be determined by the investigator.

**Positive Control:** Raji cells. Tonsil / lymph node

**Cellular Localization:** Cell membrane

**Storage and Stability:**

Ab with sodium azide is stable for 24 months when stored at 2-8°C. Antibody WITHOUT sodium azide is stable for 36 months when stored at below 0°C.

**Supplied As:**

200µg/ml of antibody purified from ascites fluid by Protein G chromatography. Prepared in 10mM PBS, pH 7.4, with 0.2% BSA and 0.09% sodium azide. Also available without BSA and azide at 1mg/ml. Or Prediluted antibody which is ready-to-use for staining of formalin-fixed, paraffin-embedded tissues.

**Key References:**

1. loachim HL; et al. International Journal of Cancer. Supplement, 1994, 8:132-3.
2. Kumar S; et al. American Journal of Hematology, 1994, 46(2):134-7.

**Limitations and Warranty:**

Our products are intended FOR RESEARCH USE ONLY and are not approved for clinical diagnosis, drug use or therapeutic procedures. No products are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our data sheets and website. Our warranty is limited to the actual price paid for the product. Lab Vision is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.

**Material Safety Data:**

This product is not licensed or approved for administration to humans or to animals other than the experimental animals. Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. Appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. The material contains 0.09% sodium azide as a preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material as indicated above. The National Institute of Occupational Safety and Health has issued a bulletin citing the potential explosion hazard due to the reaction of sodium azide with copper, lead, brass, or solder in the plumbing systems. Sodium azide forms hydrazoic acid in acidic conditions and should be discarded in a large volume of running water to avoid deposits forming in metal drainage pipes.

**For Research Use Only**

## CD74 Ab-1 (Clone LN2)

Mouse Monoclonal Antibody

**Cat. #MS-131-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml)** (Purified Ab with BSA and Azide)

**Cat. #MS-131-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml)** (Purified Ab without BSA and Azide)

**Cat. #MS-131-R7 (7.0ml)** (Ready-to-Use for Immunohistochemistry)


**Cat. #MS-131-PCS (5 Slides)** (Positive Control for Histology)

**Cat. #MS-131-PCL (0.1ml)** (Positive Control for Western Blot)

Please note this data sheet has been changed effective December 6, 2011

### Additional Key References:

1. Kanavoras P; Mikol J; Nemeth J; Galian A; Vaunaize J; Morinet F; Thurel C. Primary T-cell malignant lymphoma of the central nervous system: Histological, immunohistochemical and ultrastructural study of a case. *Path Res Pract*, 1993, 189: 93-8.
2. Perkins SL; Kjeldsberg CR. Immunophenotyping of lymphomas and leukemias in paraffin-embedded tissues. *American J of Clinical Pathology*, 1993, 99: 362-73.
3. Poston RN; Hussain IF. The immunohistochemical heterogeneity of atheroma macrophages: comparison with lymphoid tissues suggests that recently blood-derived macrophages can be distinguished from longer-resident cells. *Journal of Histochemistry and Cytochemistry*, 1993, 41:1503-12.
4. Sepp N; Radaszkiewicz T; Meijer CJ; Smolle J; Seewann H; Fritsch P; Kerl H. Specific skin manifestations in acute leukemia with monocytic differentiation. A morphologic and immunohistochemical study of 11 cases. *Cancer*, 1993, 71:124-32.
5. Shin SS; Ben-Ezra J; Burke JS; Sheibani K; Rappaport H. Reed-Sternberg-like cells in low-grade lymphomas are transformed neoplastic cells of B-cell lineage. *American Journal of Clinical Pathology*, 1993, 99(6):658-62.
6. Tallini G; West B; Buckley PJ. Diagnosis of gastrointestinal T-cell lymphomas in routinely processed tissues. *J Clin Gastroenterol*, 1993, 17: 57-66.
7. Baddoura FK; Hanson C; Chan WC. Plasmacytoid monocyte proliferation associated with myeloproliferative disorders. *Cancer*, 1992, 69(6):1457-67.
8. Carbone; Pinto A; Gloghini A; Volpe R; Zagonel V. B-zone small lymphocytic lymphoma: A morphologic, immunophenotypic, and clonality study with comparison to "well differentiated" lymphocytic disorders. *Hum Pathol*, 1992, 23: 438-48.
9. Chu WS; Abbondanzo SL; Frizzera G. Inconsistency of the immunophenotype of Reed-Sternberg cells in simultaneous and consecutive specimens from the same patients. A paraffin section evaluation in 56 patients. *Am Journal of Pathology*, 1992, 141:11-7.
10. Fukuda T; Ohnishi Y; Emura I; Tachikawa S. Microcytic variant of thymoma: histological and immunohistochemical findings in two cases. *Virchows Archiv. a, Pathological Anatomy and Histopathology*, 1992, 420(2):185-9.
11. Lavergne A; Kanavoras P; Galian A. Primary B-cell gastric lymphomas of mucosa-associated lymphoid tissue. Histological and immunohistochemical study of ten cases on surgical specimens. *Histology and Histopathology*, 1992, 7(1):129-36.
12. Lee J; Sun NC; Salahi W; Chen H; Yang MH. A clinicopathologic study on the diffuse malignant lymphoma--a morphologic and immunophenotypic analysis in 62 patients at Harbor-UCLA Medical Center. *Journal of Korean Medical Science*, 1992, 7(3):204-13.
13. O'Keane JC; Stahl D; Sheahan K; Burke B; Gottlieb LS; O'Brien MJ. Analysis of epithelial and lymphoid phenotypic markers in relation to growth pattern of colorectal adenomas. *Human Pathology*, 1992, 23(9):1038-43.
14. Ross CW; Hanson CA; Schnitzer B. CD30 (Ki-1)-positive, anaplastic large cell lymphoma mimicking gastrointestinal carcinoma. *Cancer*, 1992, 70(10):2517-23.
15. Sarker AB; Akagi T; Jeon HJ; Miyake K; Murakami I; Yoshino T; Takahashi K; Nose S. Bauhinia purpurea--a new paraffin section marker for Reed-Sternberg cells of Hodgkin's disease. A comparison with Leu-M1 (CD15), LN2 (CD74), peanut agglutinin, and Ber-H2 (CD30). *American Journal of Pathology*, 1992, 141(1):19-23.
16. Shin SS; Sheibani K; Kezirian J; Nademanee A; Forman SJ; Lee SK; Winberg CD. Immunoarchitecture of normal human bone marrow: a study of frozen and fixed tissue sections. *Human Pathology*, 1992, 23(6):686-94.
17. Stroup R; Sheibani K. Antigenic phenotypes of hairy cell leukemia and monocytoid B-cell lymphoma: an immunohistochemical evaluation of 66 cases. *Human Pathology*, 1992, 23(2):172-7.
18. Sumiyoshi Y; Kikuchi M; Takeshita M; Yoneda S; Kobari S; Ohshima K. Immunohistological study of skin involvement in Kikuchi's disease. *Virchows Archiv. B. Cell Pathology*, 1992, 62(4):263-9.
19. Takahashi H; Cheng J; Fujita S; Tsuda N; Tezuka F; Liu AR; Okabe H. Primary malignant lymphoma of the salivary gland: a tumor of mucosa-associated lymphoid tissue. *Journal of Oral Pathology and Medicine*, 1992, 21(7):318-25.



**CD74 Ab-1 (Clone LN2)**

Mouse Monoclonal Antibody

**Cat. #MS-131-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml)** (Purified Ab with BSA and Azide)

**Cat. #MS-131-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml)** (Purified Ab without BSA and Azide)

**Cat. #MS-131-R7 (7.0ml)** (Ready-to-Use for Immunohistochemistry)

**Cat. #MS-131-PCS (5 Slides)** (Positive Control for Histology)

**Cat. #MS-131-PCL (0.1ml)** (Positive Control for Western Blot)

Please note this data sheet has been changed effective December 6, 2011

20. Beljaards RC; Meijer CJ; Scheffer E; van der Valk P; Willemze R. Differential diagnosis of cutaneous large cell lymphomas using monoclonal antibodies reactive in paraffin-embedded skin biopsy specimens. *American Journal of Dermatopathology*, 1991, 13(4):342-9.
21. Catry L; Van den Oord J; Foets B; Missotten L. Morphologic and immunophenotypic heterogeneity of corneal dendritic cells. *Graefes Archive for Clinical and Experimental Ophthalmology*, 1991, 229(2):182-5.
22. Elghetany MT; Kurec AS; Schuehler K; Forbes BA; Duggan DB; Davey FR. Immunophenotyping of non-Hodgkin's lymphomas in paraffin-embedded tissue sections. A comparison with genotypic analysis. *American Journal of Clinical Pathology*, 1991, 95:517-25.
23. Gaulier A; Sabatier P; Prevot S; Fournier JG. Do measles early giant cells result from fusion of non-infected cells? An immunohistochemical and in situ hybridization study in a case of morbillous appendicitis. *Virchows Archiv. a, Pathological Anatomy and Histopathology*, 1991, 419(3):245-9.
24. Pich A; Gastaldi M; Tragni G; Navone R. Lymphocyte subsets in bone marrow lymphoid nodules and malignant lymphoma nodular involvement. *European Journal of Basic and Applied Histochemistry*, 1991, 35(1):81-9.
25. Pich A; Pisani P; Kzengli M; Cappello N; Navone R. Argyrophilic nucleolar organiser region counts and prognosis in pharyngeal carcinoma. *British Journal of Cancer*, 1991, 64:327-32.
26. Pileri S, Falini B, Sabatini E, Bigerna B, Gherlinzonii F, and Tazzari PL. Immunohistochemistry of malignant lymphomas. Advantages and limitations of the new monoclonal antibodies working in paraffin sections. *Haematologica* 76: 226-234, 1991.
27. Santucci M; Pimpinelli N; Arganini L. Primary cutaneous B-cell lymphoma: A unique type of low-grade lymphoma, Clinicopathologic and immunologic study of 83 cases. *Cancer*, 1991, 67: 2311-2326
28. Taubenberger JK; Cole DE; Raffeld M; Poplack DG; Jaffe ES; Medeiros LJ. Immunophenotypic analysis of acute lymphoblastic leukemia using routinely processed bone marrow specimens. *Archives of Pathology and Laboratory Medicine*, 1991, 115:338-42.
29. Carbone A; Pinto A; Gloghini A; De Re V; Alosi M; Zagonel V; Tirelli U; Attadia V; Boiocchi M; Volpe R. Report of an unusual small lymphocytic B-cell lymphoma selectively involving the B-zone of lymph node. *Cancer*, 1990, 66:302-12.
30. Casey TT; Cousar JB; Mangum M; Williams ME; Lee JT; Greer JP; Collins RD. Monomorphic lymphomas arising in patients with Hodgkin's disease. Correlation of morphologic, immuno-phenotypic, and molecular genetic findings in 12 cases. *American Journal of Pathology*, 1990, 136(1):81-94.
31. Cerroni L; Smolle J; Soyer P; Aparicio AM; Kerl H. Immunophenotyping of cutaneous lymphoid infiltrates in frozen and paraffin-embedded tissue sections: A comparative study. *J Am Acad Dermatol*, 1990, 22: 405-13.
32. Davey FR; Elghetany MT; Kurec AS. Immunophenotyping of hematologic neoplasms in paraffin-embedded tissue sections. *American J of Clinical Pathology*, 1990, 93(4 Suppl 1):S17-26.
33. Duggan MJ; Weisenburger DD; Ye YL; Bast MA; Pierson JL; Linder J; Armitage JO. Mantle zone lymphoma. A clinicopathologic study of 22 cases. *Cancer*, 1990, 66(3):522-9.
34. Hassan IB; Hagberg H; Sundstrom C. Immunophenotype of hairy-cell leukemia. *European Journal of Haematology*, 1990, 45(3):172-6.
35. Imam A; Stathopoulos E; Holland SL; Epstein AL; Taylor CR. Characterization of a cell surface molecule expressed on B-lymphocytes and Hodgkin's cells. *Cancer Research*, 1990, 50(5):1650-7.
36. Kanavros P; Mikol J; Nemeth J; Galian A; Dupont B; Thiebaut JB; Thurel C. Stereotactic biopsy diagnosis of primary non-Hodgkin's lymphoma of the central nervous system. A histological and immunohistochemical study. *Pathology, Research and Practice*, 1990, 186:459-66.
37. Kornstein MJ; Kay S. B cells in thymomas. *Modern Pathology*, 1990, 3(1):61-3.
38. Kurec AS; Cruz VE; Barrett D; Mason DY; Davey FR. Immunophenotyping of acute leukemias using paraffin-embedded tissue sections. *American Journal of Pathology*, 1990, 93:502-9.
39. Strickler JG; Schmidt CM; Wick MR. Methods in pathology. Immunophenotype of hairy cell leukemia in paraffin sections. *Modern Pathology*, 1990, 3(4):518-23.
40. Timens W; Kamps WA; Rozeboom-Uiterwijk T; Poppema S. Haemopoiesis in human fetal and embryonic liver. Immunohistochemical determination



## CD74 Ab-1 (Clone LN2)

### Mouse Monoclonal Antibody

**Cat. #MS-131-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml)** (Purified Ab with BSA and Azide)

**Cat. #MS-131-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml)** (Purified Ab without BSA and Azide)


**Cat. #MS-131-R7 (7.0ml)** (Ready-to-Use for Immunohistochemistry)

**Cat. #MS-131-PCS (5 Slides)** (Positive Control for Histology)

**Cat. #MS-131-PCL (0.1ml)** (Positive Control for Western Blot)

Please note this data sheet has been changed effective December 6, 2011

- in B5-fixed paraffin-embedded tissues. Virchows Archiv. a, Pathological Anatomy and Histopathology, 1990, 416(5):429-36.
41. Valente G; Ferrara P; Stramignoni A. Lymphocyte populations of non-sclerodular Hodgkin's disease subtypes in different stages of lymphocyte depletion. An immunophenotypic and quantitative study. Virchows Archiv. B. Cell Pathology, 1990, 58(4):289-94.
  42. van den Oord JJ; De Vos R; Facchetti F; Delabie J; De Wolf-Peeters C; Desmet VJ. Distribution of non-lymphoid, inflammatory cells in chronic HBV infection. Journal of Pathology, 1990, 160:223-30.
  43. Wick MR; Mills SE; Swanson PE. Expression of "myelomonocytic" antigens in mesotheliomas and adenocarcinomas involving the serosal surfaces. American Journal of Clinical Pathology, 1990, 94(1):18-26.
  44. Yoshino T; Hoshida Y; Murakami I; Takahashi K; Akagi T. Comparison of monoclonal antibodies reactive with lymphocyte subsets in routinely fixed paraffin-embedded material: flow cytometric analyses, immunoperoxidase staining and influence of fixatives. Acta Medica Okayama, 1990, 44(5):243-50.
  45. Carbone A; Gloghini A; Pinto A; Attadia V; Zagonel V; Volpe R. Monocytoid B-cell lymphoma with bone marrow and peripheral blood involvement at presentation [see comments]. American Journal of Clinical Pathology, 1989, 92(2):228-36.
  46. Eckert F; Schmid U. Identification of plasmacytoid T cells in lymphoid hyperplasia of the skin. Archives of Dermatology, 1989, 125:1518-24.
  47. Katzin WE; Linden MD; Fishleder AJ; Tubbs RR. Immunophenotypic and genotypic characterization of diffuse mixed non-Hodgkin's lymphomas. American Journal of Pathology, 1989, 135(4):615-21.
  48. Kubic VL; Brunning RD. Immunohistochemical evaluation of neoplasms in bone marrow biopsies using monoclonal antibodies reactive in paraffin-embedded tissue. Modern Pathol, 1989, 2:618-29.
  49. Leong AS; Gilham PN. The effects of progressive formaldehyde fixation on the preservation of tissue antigens. Pathology, 1989, 21:266-8.
  50. Manivel JC; Wick MR; Coffin CM; Dehner LP. Immunohistochemistry in the differential diagnosis in the second-look operation for ovarian carcinomas. International Journal of Gynecological Pathology, 1989, 8(2):103-13.
  51. Nakhleh RE; Manivel JC; Hurd D; Sung JH. Central nervous system lymphomas. Immunohistochemical and clinicopathologic study of 26 autopsy cases. Archives of Pathology and Laboratory Medicine, 1989, 113(9):1050-6.
  52. Norton AJ; Isaacson PG. Lymphoma phenotyping in formalin-fixed and paraffin wax-embedded tissues. I. Range of antibodies and staining patterns. Histopathol, 1989, 14:437-46.
  53. O'Brian DS; Kennedy MJ; Daly PA; O'Brien AA; Tanner WA; Rogers P; Lawlor E. Multiple lymphomatous polyposis of the gastrointestinal tract. A clinicopathologically distinctive form of non-Hodgkin's lymphoma of B-cell centrocytic type. American Journal of Surgical Pathology, 1989, 13(8):691-9.
  54. Wick MR; Mills SE; Dehner LP; Bollinger DJ; Fechner RE. Serous papillary carcinomas arising from the peritoneum and ovaries. A clinicopathologic and immunohistochemical comparison. International Journal of Gynecological Pathology, 1989, 8:179-88.
  55. Andrade RE; Wick MR; Frizzera G; Gajl-Peczalska KJ. Immunophenotyping of hematopoietic malignancies in paraffin sections. Human Pathology, 1988, 19(4):394-402.
  56. Ashby MA; Barber PC; Holmes AE; Freer CE; Collins RD. Primary intracranial Hodgkin's disease. A case report and discussion. American Journal of Surgical Pathology, 1988, 12(4):294-9.
  57. Coles FB; Cartun RW; Pastuszak WT. Hodgkin's disease, lymphocyte-predominant type: immunoreactivity with B-cell antibodies. Modern Pathology, 1988, 1(4):274-8.
  58. Davey FR; Olson S; Kurec AS; Eastman-Abaya R; Gottlieb AJ; Mason DY. The immunophenotyping of extramedullary myeloid cell tumors in paraffin-embedded tissue sections. American Journal of Surgical Pathology, 1988, 12(9):699-707.
  59. Facchetti F; De Wolf-Peeters C; van den Oord JJ; De vos R; Desmet VJ. Plasmacytoid T cells: a cell population normally present in the reactive lymph node. An immunohistochemical and electronmicroscopic study. Human Pathology, 1988, 19:1085-92.
  60. Facchetti F; De Wolf-Peeters C; Van den Oord JJ; Desmet VJ. Plasmacytoid T cells in a case of lymphocytic infiltration of skin. A component of the skin-associated lymphoid tissue? Journal of Pathology, 1988, 155(4):295-300.



**CD74 Ab-1 (Clone LN2)**

## Mouse Monoclonal Antibody

**Cat. #MS-131-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml)** (Purified Ab with BSA and Azide)**Cat. #MS-131-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml)** (Purified Ab without BSA and Azide)**Cat. #MS-131-R7 (7.0ml)** (Ready-to-Use for Immunohistochemistry)**Cat. #MS-131-PCS (5 Slides)** (Positive Control for Histology)**Cat. #MS-131-PCL (0.1ml)** (Positive Control for Western Blot)**Please note this data sheet has been changed effective December 6, 2011**

61. Linder J; Ye Y; Armitage JO; Weisenburger DD. Monoclonal antibodies marking B-cell non-Hodgkin's lymphoma in paraffin-embedded tissue. *Modern Pathology*, 1988, 1:29-34.
62. Timens W; Boes A; Rozeboom-Uiterwijk T; Poppema S. Immuno-architecture of human fetal lymphoid tissues. *Virchows Archiv. a, Pathological Anatomy and Histopathology*, 1988, 413:563-71.
63. Wieczorek R; Buck D; Bindl J; Knowles DM. Monoclonal antibody Leu-22 (L60) permits the demonstration of some neoplastic T cells in routinely fixed and paraffin-embedded tissue sections. *Human Pathology*, 1988, 19(12):1434-43.
64. Linder J; Ye Y; Harrington DS; Armitage JO; Weisenburger DD. Monoclonal antibodies marking T lymphocytes in paraffin-embedded tissue. *Am J Pathol*, 1987, 127: 1-8.
65. Ng CS; Chan JK; Lo ST; Lo DS. Critical assessment of four monoclonal antibodies reactive with B-cells in formalin-fixed paraffin-embedded tissues. *Histopathology*, 1987, 11:1243-58.
66. Norton AJ; Isaacson PG. Detailed phenotypic analysis of B-cell lymphoma using a panel of antibodies reactive in routinely fixed wax-embedded tissue. *American Journal of Pathology*, 1987, 128:225-40.
67. Norton AJ; Isaacson PG. The diagnosis of malignant lymphoma using monoclonal antibodies reactive in routinely fixed wax embedded tissue. *J of Pathol*, 1987, 151: 183-4.
68. Poppema S, Hollema H, Visser L, and Vos H. Monoclonal antibodies (MT1, MT2, MB1, MB2, MB3) reactive with leukocyte subsets in paraffin-embedded tissue sections. *Am J Pathol*, 1987, 127: 418-29.
69. Sheibani K; Battifora H; Winberg CD; Burke JS; Ben-Ezra J; Ellinger GM; Quigley NJ; Fernandez BB; Morrow D; Rappaport H. Further evidence that "malignant angioendotheliomatosis" is an angiotropic large-cell lymphoma. *New England Journal of Medicine*, 1986, 314(15):943-8.
70. Sherrod AE; Felder B; Levy N; Epstein A; Marder R; Lukes RJ; Taylor CR. Immunohistologic identification of phenotypic antigens associated with Hodgkin and Reed-sternberg cells: A paraffin section study. *Cancer*, 1986, 57: 2135-40.
71. Okhon E; Felder B; Epsteim A; Lukes RJ; Taylor CR. Monoclonal antibodies rective with B-lymphocytes and histiocytes in paraffin sections. *Cancer*, 1985, 56: 95-104.
72. Sausville EA; Worsham GF; Matthews MJ; Makuch RW; Fischmann AB; Schechter GP; Gazdar AF; Bunn PA Jr. Histologic assessment of lymph nodes in mycosis fungoides/Sezary syndrome (cutaneous T-cell lymphoma): clinical correlations and prognostic importance of a new classification system. *Human Pathology*, 1985, 16:1098-109.
73. Marder RJ, Varakojis D, Silver J, and Epstein AL. Immunohistochemical analysis of human lymphomas with monoclonal antibodies to B Cell and Ia antigens reactive in paraffin sections. *Lab Invest* 52: 497-504, 1985.
74. Epstein AL, Marder RJ, Winter JN, and Fox RI. Two new monoclonal antibodies (LN-1, LN-2) reactive in B5 formalin-fixed, paraffin-embedded tissues with follicular center and mantle zone human B lymphocytes and derived tumors. *J of Immunology* 133: 1028-1036, 1984.

