

Keratin 19 Ab-4 (Clone BA17)

Mouse Monoclonal Antibody

Cat. #MS-1671-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Purified Ab with BSA and Azide)

Cat. #MS-1671-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml) (Purified Ab without BSA and Azide)

Cat. #MS-1671-R7 (7.0ml) (Ready-to-Use for Immunohistochemical Staining)

Cat. #MS-1671-PCS (5 Slides) (Positive Control for Histology)

Cat. #MS-1671-PCL (0.1ml) (Positive Control for Western Blot)

Description: Cytokeratin 19 is a member of type I acidic subfamily of intermediate filaments. It is expressed in various different human tissues except in liver. Keratin 19 is not expressed in hepatocytes, therefore, antibody to keratin 19 is useful in the identification of liver metastasis.

Mol. Wt. of Antigen: 40kDa

Epitope: Not determined

Species Reactivity: Human. Others-not known.

Clone Designation:BA17

Ig Isotype / Light Chain: IgG₁ / κ

Immunogen: Detergent soluble extract of human mammary epithelium

Applications and Suggested Dilutions:

- Western Blotting (Ab 2-4µg/ml for 2 hrs at RT)
- Immunohistology (Formalin/paraffin) (Ab 1:200 – 1:400 for 30 min at RT)
- * [Staining of formalin-fixed tissues REQUIRES boiling tissue sections in 10mM citrate buffer, pH 6.0, (**NEOMARKERS'** Cat. #AP-9003), for 10-20 min followed by cooling at RT for 20 min.]

The optimal dilution for a specific application should be determined by the investigator.

Positive Control: Ls174T and HT29 Cells, Colon ca

Cellular Localization: Cytoplasmic

Supplied As:

200µg/ml antibody purified from the ascites fluid by Protein G chromatography. Prepared in 10mM PBS, pH 7.4, with 0.2% BSA and 0.09% sodium azide. Also available without BSA and azide at 1mg/ml.

or

Prediluted antibody which is ready-to-use for staining of formalin-fixed, paraffin-embedded tissues.

Storage and Stability: Ab with sodium azide is stable for 24 months when stored at 2-8°C. Antibody WITHOUT sodium azide is stable for 36 months when stored at below 0°C.

Key References:

1. Bartek J, et al. (1985) J Cell Sci, 75:17-33.
2. Moll R, et al. (1982) Cell, 31:11-24.
3. Tseng S C G, et al. (1982) Cell, 30:361-72.
4. Bartek J, et al. (1986) Histochem J, 18:565-575.
5. Bartek J et al. (1985) Int J Cancer, 36:299-306.
6. Bartek J et al. (1986) Eur J Cancer Clin Oncol, 22:1441-52.
7. Bartkova J, et al. (1987) Tumor Biol, 8:45-56.

Limitations and Warranty:

Our products are intended FOR RESEARCH USE ONLY and are not approved for clinical diagnosis, drug use or therapeutic procedures. No products are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our data sheets and website. Our warranty is limited to the actual price paid for the product. NeoMarkers is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.

Material Safety Data:

This product is not licensed or approved for administration to humans or to animals other than the experimental animals. Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. Appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. The material contains 0.09% sodium azide as a preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material as indicated above. The National Institute of Occupational Safety and Health has issued a bulletin citing the potential explosion hazard due to the reaction of sodium azide with copper, lead, brass, or solder in the plumbing systems. Sodium azide forms hydrazoic acid in acidic conditions and should be discarded in a large volume of running water to avoid deposits forming in metal drainage pipes.

For Research Use Only

Suggested References:



Keratin 19 Ab-4 (Clone BA17)

Mouse Monoclonal Antibody

Cat. #MS-1671-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Purified Ab with BSA and Azide)

Cat. #MS-1671-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml) (Purified Ab without BSA and Azide)

Cat. #MS-1671-R7 (7.0ml) (Ready-to-Use for Immunohistochemical Staining)

Cat. #MS-1671-PCS (5 Slides) (Positive Control for Histology)

Cat. #MS-1671-PCL (0.1ml) (Positive Control for Western Blot)

- Hazelbag HM; van den Broek LJ; van Dorst EB; Offerhaus GJ; Fleuren GJ; Hogendoorn PC. Immunostaining of chain-specific keratins on formalin-fixed, paraffin-embedded tissues: a comparison of various antigen retrieval systems using microwave heating and proteolytic pre-treatments. *Journal of Histochemistry and Cytochemistry*, 1995, 43(4):429-37.
- Narisawa Y; Hashimoto K; Kohda H. Immunohistochemical demonstration of keratin 19 expression in isolated human hair follicles. *Journal of Investigative Dermatology*, 1994, 103(2):191-5.
- Akasofu M; Kawahara E; Kurumaya H; Nakanishi I. Immunohistochemical detection of breast specific antigens and cytokeratins in metastatic breast carcinoma in the liver. *Acta Pathologica Japonica*, 1993, 43(12):736-44.
- Klijanienko J; el-Naggar A; De Braud F; Micheau C; Janot F; Luboinski B; Gentile A; Russo A; Cvitkovic E. Keratins 6, 13 and 19. Differential expression in squamous cell carcinoma of the head and neck. *Analytical and Quantitative Cytology and Histology*, 1993, 15(5):335-40.
- Nasser IA; Lee AK; Bosari S; Saganich R; Heatley G; Silverman ML. Occult axillary lymph node metastases in "node-negative" breast carcinoma. *Human Pathology*, 1993, 24(9):950-7.
- Su L; Morgan PR; Thomas JA; Lane EB. Expression of keratin 14 and 19 mRNA and protein in normal oral epithelia, hairy leukoplakia, tongue biting and white sponge nevus. *Journal of Oral Pathology and Medicine*, 1993, 22(4):183-9.
- Coltrera MD; Zarbo RJ; Sakr WA; Gown AM. Markers for dysplasia of the upper aerodigestive tract. Suprabasal expression of PCNA, p53, and CK19 in alcohol-fixed, embedded tissue. *American Journal of Pathology*, 1992, 141(4):817-25.
- Mittal KR; Demopoulos RI; Goswami S. Patterns of keratin 19 expression in normal, metaplastic, condylomatous, atrophic, dysplastic, and malignant cervical squamous epithelium. *American Journal of Clinical Pathology*, 1992, 98(4):419-23.
- Nomoto M; Uchikosi Y; Kajikazawa N; Tanaka Y; Asakura H. Appearance of hepatocytelike cells in the interlobular bile ducts of human liver in various liver disease states. *Hepatology*, 1992, 16:1199-205.
- Carmichael RP; McCulloch CA; Zarb GA. Immunohistochemical localization and quantification of desmoplakins I & II and keratins 1 and 19 in plastic-embedded sections of human gingiva. *Journal of Histochemistry and Cytochemistry*, 1991, 39(4):519-28.
- Bader BL; Franke WW. Cell type-specific and efficient synthesis of human cytokeratin 19 in transgenic mice. *Differentiation*, 1990, 45:109-18.
- Bartek J; Bartkova J; Taylor-Papadimitriou J. Keratin 19 expression in the adult and developing human mammary gland. *Histochemical Journal*, 1990, 22(10):537-44.
- Leoncini P; Petracca R; Ruggiero P; Cintorino M; Syrjanen S; Mantyljarvi R; Syrjanen K. Expression of cytokeratin No. 19 polypeptide in genital papillomavirus lesions. *Gynecologic and Obstetric Investigation*, 1990, 29(1):59-66.
- Stosiek P; Kasper M; Karsten U. Expression of cytokeratin 19 during human liver organogenesis. *Liver*, 1990, 10(1):59-63.
- Van Eyken P; Sciot R; Callea F; Ramaekers F; Schaart G; Desmet VJ. A cytokeratin-immunohistochemical study of hepatoblastoma. *Human Pathology*, 1990, 21(3):302-8.
- Van Eyken P; Sciot R; Callea F; Ramaekers F; Schaart G; Desmet VJ. A cytokeratin-immunohistochemical study of hepatoblastoma. *Human Pathology*, 1990, 21(3):302-8.

