

Ku (p70) Ab-4 (Clone N3H10)

Mouse Monoclonal Antibody

Cat. #MS-329-P0, -P1, or -P (0.1ml, 0.5ml, or 1.0ml at 200µg/ml) (Purified with BSA and Azide)

Cat. #MS-329-P1ABX or -PABX (0.1ml or 0.2ml at 1.0mg/ml) (Purified without BSA and Azide)

Cat. #MS-329-R7 (7.0ml) (Ready-to-Use for Immunohistochemical Staining)

Cat. #MS-329-PCS (5 Slides) (Positive Control for Histology)

Cat. #MS-329-PCL (0.1ml) (Positive Control for Western Blot)

Please note this data sheet has been changed effective December 6, 2011

Description: The Ku autoantigen is a heterodimer of 70kDa (p70) and ~80kDa (p80) proteins. The p70/p80 dimer is important for function of a 460kDa DNA-dependent protein kinase that phosphorylates certain transcription factors, including Sp1, Oct-1, p53, and SV40 large T antigen *in vitro*. Ku protein plays a role in cell signaling, proliferation, DNA repair, replication, transcriptional activation, and apoptosis.

Mol. Wt. of Antigen: 70kDa

Epitope: aa 506-541

Species Reactivity: Human, Monkey, and *Xenopus*. Mouse & Hamster (Western blotting only). Does not react with cow, rabbit, and sea urchin.

Clone Designation: N3H10

Ig Isotype: IgG_{2b}

Immunogen: Human placental extract designated as PSE1-PL

Applications and Suggested Dilutions:

- Immunofluorescence
- Immunoprecipitation (Native and denatured) (Use Protein A) (Ab 2µg/mg protein lysate)
- Western Blotting (0.25-0.5µg/ml for 2hrs at RT)
- Immunohistology (Formalinparaffin) (1:200-400 for 20 minutes at RT using the LP system)

* [Staining of formalin-fixed tissues is IMPROVED by boiling tissue sections in 10mM citrate buffer, pH 6.0, (**NEOMARKERS'** Cat. #AP-9003), for 10-20 min followed by cooling at RT for 20 min.]

The optimal dilution for a specific application should be determined by the investigator.

Positive Control: BT474, T47D, HEP-G-2 or HeLa cells. Tonsil

Cellular Localization: Nuclear

Storage and Stability: Ab with sodium azide is stable for 24 months when stored at 2-8°C. Antibody WITHOUT sodium azide is stable for 36 months when stored at below 0°C.

Supplied As: 200µg/ml antibody purified from the ascites fluid by Protein A chromatography. Prepared in 10mM PBS, pH 7.4, with 0.2% BSA and 0.09% sodium azide. Also available without BSA and azide at 1mg/ml. **Or** Prediluted antibody which is ready-to-use for staining of formalin-fixed, paraffin-embedded tissues.

Key References:

1. Ajmani AK, *et. al.* J Exp Med, 1995, 181:2049-58.
2. Wang J, *et. al.* Mol Biol Reports, 1993, 18:15-28.

Limitations and Warranty:

Our products are intended FOR RESEARCH USE ONLY and are not approved for clinical diagnosis, drug use or therapeutic procedures. No products are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our data sheets and website. Our warranty is limited to the actual price paid for the product. NeoMarkers is not liable for any property damage, personal injury, time or effort or economic loss caused by our products.


Material Safety Data:

This product is not licensed or approved for administration to humans or to animals other than the experimental animals. Standard Laboratory Practices should be followed when handling this material. The chemical, physical, and toxicological properties of this material have not been thoroughly investigated. Appropriate measures should be taken to avoid skin and eye contact, inhalation, and ingestion. The material contains 0.09% sodium azide as a preservative. Although the quantity of azide is very small, appropriate care should be taken when handling this material as indicated above. The National Institute of Occupational Safety and Health has issued a bulletin citing the potential explosion hazard due to the reaction of sodium azide with copper, lead, brass, or solder in the plumbing systems. Sodium azide forms hydrazoic acid in acidic conditions and should be discarded in a large volume of running water to avoid deposits forming in metal drainage pipes.

For Research Use Only



Lab Vision Corporation
46360 Fremont Blvd.
Fremont, CA 94538-6406, USA
US Toll Free: 1 (800) 522-7270
Phone: +1 (269) 544-5600
Fax: 1 (269) 372-2674
www.thermoscientific.com/labvision



Thermo Fisher Scientific
Anatomical Pathology
Tudor Road, Manor Park
Runcorn, Cheshire WA7 1TA, UK
Tel: +44 (0) 1928 534 050
Fax: +44 (0) 1928 534 049
sales.ap.uk@thermofisher.com