



SPECIFICATION SHEET

Hydrogen Peroxide Block

DESCRIPTION:

Hydrogen Peroxide Block has been formulated for use with immunohistochemical techniques. It is designed to quench endogenous peroxidase that can cause nonspecific background staining in enzyme labeling methods. Peroxidase within red blood cells may be difficult to quench entirely.

AVAILABILITY:

<u>CATALOG #</u>	<u>VOLUME</u>
TA-060-HP	60ml
TA-125-HP	125ml

IMMUNOHISTOCHEMICAL STAINING PROTOCOL:

- Hydrogen Peroxide Block may be applied after the sections have been deparaffinized and rehydrated. The application must occur prior to the primary antibody incubation. Incubate tissue section for 10-15 minutes at room temperature or 37° C. After incubation, rinse twice in buffer.
- For bulk staining, pour Hydrogen Peroxide Block in a covered staining tray and dip slides for 5-10 minutes. Replace with fresh Hydrogen Peroxide Block after 5-10 uses. This step can be performed at the time of deparaffinization.

STORAGE:


Store at 2-8°C. Product is stable for 18 months from date of manufacture.

PRECAUTIONS:

47790 Westinghouse Dr. Fremont CA 94539 USA
Tel: (800) 828-1628 Fax: (510) 991-2826
Email: labvision@labvision.com Website: www.labvision.com



TA-060-HP
Rev 21204
Page 1 Of 1



Do not pipette reagent by mouth. Avoid contact with skin and eyes. Wash after use. Observe all federal, state and local environmental regulations regarding disposal.

LIMITATIONS:

This product is available for research use only; not for diagnostic or therapeutic work. Lab Vision Corporation will not be held responsible for patent infringement or other violation that may occur with the use of this product.